





Naming Hydrocarbons (nomenclature)



Drawing Structures: It's All Good



This is called the
"condensed structure"



On a test, choose a
method that shows all Hs

Using brackets can also shorten some formulas:



Basic Naming of Hydrocarbons

Hydrocarbon names are based on: 1) type,
2) # of carbons, 3) side chain type and position

1) name will end in -ane, -ene, or -yne

2) the number of carbons is given by a "prefix"

1 meth- 2 eth- 3 prop- 4 but- 5 pent-

6 hex- 7 hept- 8 oct- 9 non- 10 dec-

Actually, all end in a, but a is dropped when next
to a vowel. E.g. a 6 C alkene is hexene

Q - What names would be given to these:

7C, 9C alkane


2C, 4C alkyne

1C, 3C alkene

Numbering Carbons


Q- draw pentene

- Naming compounds with multiple bonds is more complex than previously indicated.
- When 2+ possibilities exist, #s are needed.
- Always give double bond the lowest number.
- Q - Name these



Naming Side Chains

- The names of molecules with branches are based on: side chains, root
2,3-dimethylpentane




- The "root" or "parent chain" is usually the longest possible hydrocarbon chain.
- The root must include multiple bonds if they are present. If a cyclic structure is present it will be the root even if it is not the longest chain.
- Side chains are also called "side branches" or "alkyl groups". Their names end in -yl.

Common side chains :

-CH₃ methyl, -CH₂CH₃ ethyl, -CH₂CH₂CH₃ propyl

Naming Side Chains

Example: use the rules on this handout
to name the following structure



IUPAC Rules for Naming Hydrocarbons

- Choose the correct ending: -ane, -ene, or -yne
- Determine the longest carbon chain. Where a double or triple bond is present, choose the longest chain that includes this bond. If there is a cyclic structure present, the longest chain starts and stops within the cyclic structure.
- Assign numbers to each C of the parent chain. For alkenes and alkynes the first carbon of the multiple bond should have the smallest number. For alkanes the first branch (or first point of difference) should have the lowest #. Carbons in a multiple bond must be numbered consecutively.
- Attach a prefix that corresponds to the number of carbons in the parent chain. Add cyclo- to the prefix if it is a cyclic structure.

IUPAC Rules for Naming Hydrocarbons

- Determine the correct name for each branch ("alkyl" groups include methyl, ethyl, propyl, etc.)
- Attach the name of the branches alphabetically, along with their carbon position, to the front of the parent chain name. Separate numbers from letters with hyphens (e.g. 4-ethyl-2-methyldecane)
- When two or more branches are identical, use prefixes (di-, tri-, etc.) (e.g. 2,4-dimethylhexane). Numbers are separated with commas. Prefixes are ignored when determining alphabetical order. (e.g. 2,3,5-trimethyl-4-propylheptane)
- When identical groups are on the same carbon, repeat the number of this carbon in the name. (e.g. 2,2-dimethylhexane)